## ESTIMATION OF MONTHLY REFERENCE EVAPOTRANSPIRATION WITH SCARCE INFORMATION USING MACHINE LEARNING IN SOUTHWESTERN COLOMBIA

Meteoro

Juan Camilo Triana-Madrid,<sup>1</sup> Camilo Ocampo-Marulanda<sup>2,3</sup>, Yesid Carvajal-Escobar<sup>4</sup> Wilmar Alexander Torres-López<sup>5</sup>, Joshua Triana<sup>6</sup> y Teresita Canchala<sup>7</sup>

<sup>1,2,4,6,7</sup>Water Resources Engineering and Soil Research Group (IREHISA), School of Natural Resources and Environmental Engineering, Universidad del Valle, Cali 25360, Colombia

<sup>3</sup>Faculty of Natural Sciences and Engineering, Fundación Universitaria de San Gil, Yopal 850007, Colombia

<sup>5</sup>Research Group in Applied Statistics (INFERIR), School of Statistics, Universidad del Valle, Cali 25360, Colombia

<sup>6</sup>Univalle Group in Artificial Intelligence (GUIA), School of Systems and Computer Engineering, Universidad del Valle, Cali 25360, Colombia

Autor correspondiente. Yesid Carvajal, vesid.carvajal@correounivalle.edu.co

Manuscrito recibido el 11 de octubre de 2022, en su versión final el 16 de diciembre de 2022

#### ABSTRACT

This research aimed to identify an alternative method to estimate reference evapotranspiration (ETo) with scarce climatological information in southwestern Colombia between 1983-2017 by evaluating and comparing different machine learning techniques. The FAO Penman-Monteith (*FAO-PM56*) was used as the reference method and four empirical methods (*Hargreaves, Thornthwaite, Cenicafé, and Turc*) were assessed with five metrics to



evaluate the method of best fit to *FAO-PM56*, root mean square error (RMSE), mean absolute error (MAE), mean bias error (MBE), Nash-Sutcliffe model efficiency coefficient (NSE), and Pearson correlation coefficient (R). Three models were designed using machine learning techniques to estimate ETo, multiple linear regression (MLR), artificial neural networks (ANN), and autoregressive integrated moving average model (ARIMA). The results showed that the ARIMA-M3 model reported the best performance metrics (RMSE = 4.13 mm month<sup>-1</sup>, MAE = 3.15 mm month<sup>-1</sup>, MBE = -0.08 mm month<sup>-1</sup>, NSE = 0.96 and r = 0.98). However, it restricts in that it can only be used locally and cannot be extrapolated to other climatological stations, because it was calibrated with specific conditions (exogenous variables) and stations, unlike the ANN-M1 model, which only requires training the network for its application. This method will allow estimating ETo in places with scarce information, as vital for water management in places with much uncertainty regarding accessibility and availability.

**Key Words:** Artificial neural network; FAO-56 Penman-Monteith; Performance metrics; Southwestern Colombia; Evapotranspiration

## ESTIMACIÓN DE EVAPOTRANSPIRACIÓN DE REFERENCIA CON INFORMACIÓN ESCASA UTILIZANDO MACHINE LEARNING EN EL SUROCCIDENTE COLOMBIANO

#### RESUMEN

Esta investigación tuvo como objetivo identificar un método alternativo para estimar la evapotranspiración de referencia (ETo) con escasa información climatológica en el suroeste de Colombia entre 1983-2017, evaluando y comparando diferentes técnicas de machine learning. Se utilizó el método de FAO Penman-Monteith (*FAO-PM56*) como método de referencia y se evaluaron 4 métodos de empíricos (*Hargreaves, Thornthwaite, Cenicafé y Turc*) con cinco métricas para evaluar el método de mejor ajuste al FAO-PM56, error



cuadrático medio (RMSE), error medio absoluto (MAE), error medio de sesgo (MBE), coeficiente de eficiencia del modelo de Nash-Sutcliffe (NSE) y coeficiente de correlación de Pearson (R). Se diseñaron tres modelos utilizando técnicas de machine learning para estimar la ETo, regresión lineal múltiple (MLR), redes neuronales artificiales (ANN) y modelo de media móvil integrada autorregresiva (ARIMA). Los resultados mostraron que el modelo ARIMA-M3 presentó la mejor métrica de rendimiento (RMSE = 4,13 mm mes<sup>-1</sup>, MAE = 3,15 mm mes<sup>-1</sup>, MBE = -0,08 mm mes<sup>-1</sup>, NSE = 0,96 y R = 0,98). Sin embargo, tiene la restricción de que sólo se puede utilizar localmente y no se puede extrapolar a otras estaciones climatológicas, porque se calibró con estaciones y condiciones específicas (variables exógenas), a diferencia del modelo RNA-M1, que sólo requiere entrenar la red para su aplicación. Este método permitirá estimar la ETo en lugares con escasa información, lo que es vital para la gestión del agua en lugares con mucha incertidumbre en cuanto a accesibilidad y disponibilidad.

Palabras clave: Redes neuronales artificiales; FAO-PM56 Penman-Monteith, Métricas de desempeño, Suroccidente Colombiano, Evapotranspiración

#### 1) INTRODUCTION

Evapotranspiration is an aerodynamic physical process where water from the land surface evaporates, and water from plants transpires (Allen et al., 1998; Alves et al., 2017; Granata et al., 2020; Meneses et al., 2020). Correct estimation of evapotranspiration is fundamental in different research areas (Maček et al. 2018; Kumar et al., 2020), such as climate change (Cannarozo et al., 2006; Liu et al., 2008; Yao et al. 2009; Yang et al., 2011), hydroclimatology (Rivas and Caselles 2004; Castañeda and Rao, 2005), water resources planning and management (Huizhi and Jianwu, 2012; Łabędzki et al., 2014; Biggs et al., 2016) and irrigation needs (Yoder et al., 2005; Tabari, 2010).

The lysimeter is a suitable method for estimating field evapotranspiration in the field (Wang and Dickinson 2012). However, it has several limitations, such as high installation costs, complex instruments (Valipour, 2015; Goh et al., 2021), prior experimental setup, and maintenance hours to achieve reliable results (Igbadun et al., 2006; Choi and Jeon 2018; Jing et al., 2019; Ahmadi and Javanbakht 2020).

Indirect methods have been developed to estimate reference evapotranspiration (ETo) (Choi and Jeon, 2018). ETo is defined as the evapotranspiration of a grass crop with an assumed height of 0.12 m, a fixed surface resistance of 70 s m<sup>-1</sup>, and an albedo of 0.23, with adequate nutrient and water availability (Allen et al., 1998). Methods that estimate ETo based on climatic factors can be classified as those based on temperature (Thornthwaite and Wilm, 1948; Blanney and Criddle, 1950; Hargreaves and Samani, 1985), radiation (Turc, 1961; Priestley and Taylor 1972), and combined methods (Allen et al., 1998).

The most recommended method for ETo estimation is the so-called FAO Penman-Monteith (*FAO-PM56*), developed by the Food and Agriculture Organization of the United Nations (FAO) (Allen et al., 1998) and the World Meteorological Organization (WMO) (Allen et al., 1998; Granata et al., 2020; Cobaner, 2010; Huo et al., 2012; Laqui et al., 2014; Ayaz et al., 2021). It can be used anywhere in the world without the need to calibrate the equation. Because of this, it has been subject to extensive validation with lysimeters in various global climatic conditions (Landeras et al., 2008; Nema et al., 2017; Quej et al., 2019; Ayaz et al., 2021; Ferreira et al., 2021).

The main drawback of using this method is the requirement of a significant number of variables for its estimation, these variables are maximum and minimum air temperature, wind speed, precipitation, and solar radiation (Valipour, 2015). This restricts its worldwide use, especially in places with a lack and insufficient availability of climatological information. For example, in protected areas and/or armed conflict countries with low budgets (Traore et al., 2008).

Numerous studies evaluate ETo models in Colombia, standing out among them the one by Barco et al. (2020), who made a macroscale estimate of evaporation in Colombia using the methods of *Turc*, *Morton, Penman, Holdridge, and Budyko*. However, they did not make comparisons with field measurements. Jaramillo (2006) developed the empirical equation of the National Coffee Research Center (Cenicafé) in several locations in the Colombian Andes, mainly in the Cauca and Magdalena River basins, and compared the relationship between the observed values of *FAO-PM56* with the Class A evaporation tank. Poveda et al. (2007) regionalized ETo in Colombia using the methods of *Turc, Morton, Coutagne, Thornthwaite, Holdridge, Meyer, Penman, Budyko, and Cenicafé*. Ramírez et al. (2011) assessed the application of the *FAO-PM56*, the Hargreaves, the Garcia and Lopez modified, and the lysimeter to estimate ETo in the coffee zone of Colombia. Toro-Trujillo et al. (2015) evaluated the reliability of evapotranspiration estimation of the *Hargreaves-Samani* and radiation methods concerning the *FAO-PM56* method in the northern banana-growing zone Urabá Antioqueño. Mendoza & Peña (2021) compared the *Blaney-Criddle, Hargreaves, Priestley-Taylor*,

*and Camargo* methods with values of class A evaporation tanks from Colombian Sugarcane Research Center (Cenicaña) and *FAO-PM56*.

Meteoro

logica

The different results of the authors agree that no method shows significant superiority over the other, which is attributed to the low quality of the available information. For this reason, models to estimate ETo have been developed during the last decades, using machine learning techniques such as Artificial Neural Networks (ANN) (Zanneti et al., 2008; Alves et al., 2017; Fonseca et al., 2018; Laqui et al., 2019; Meneses et al., 2020) Autoregressive Integrated Moving Average Model (ARIMA) (Jordan et al., 2008; Gautam and Sinha, Mossad and Alazba, 2016; Bouznad et al., 2020), and Multiple Linear Regression (MLR) (Yirga, 2019).

In Colombia, the spatial distribution of climatological stations is uneven due to several factors, such as complex topography (e.g., the Andean Mountain range), areas affected by the armed conflict, and low investment in technological resources, among others (Urrea et al., 2019; Canchala et al., 2022). In southwestern Colombia (Nariño), 76% of the rainfall stations are in the Andean region, with a density of one station every 470 km<sup>2</sup>, covering 40% of the total area of Nariño. The remaining 24% of the rainfall stations are in the Pacific region, with a density of one station every 1,442 km<sup>2</sup>, accounting the 52% of the total area of Nariño. In the case of climatological stations, the scenario is worse since, in the Andean region, there is one every 1,720 km<sup>2</sup> and only one in the Pacific region (Barbacoas) (Ocampo-Marulanda et al., 2022). The rainfall stations are those available Instituto de Hidrología, Meteorología y Estudios Ambientales (IDEAM). The accessibility and availability of information on climatological variables allow for a better understanding of the hydrological cycle and more efficient management for its use in agriculture.

The southwest of Colombia has the highest percentage of harvested area (7.9%); therefore, considering the problems mentioned earlier and that Nariño is one of the departments with the highest participation in the country's agricultural production (Moncayo, 2015), adequate planning of water resources must be carried out to ensure food security.

In this scenario, the objective of this research was to determine a model that allows estimating ETo in a scenario with scarce information and high spatiotemporal variability in climatic elements. A contradictory aspect of the study area is that the areas with higher rainfall have less climatological information and more missing data in their records. Knowledge of ETo would enable better management to contribute to the sustainability of food sovereignty and security by meeting the Sustainable Development Goals related to objective 2, zero hunger; objective 13, climate action; and objective 15, life on land (United Nations, 2018).

#### 2) DATA AND METHODS

#### 2.1) Study area

The southwest of Colombia (Nariño) is one of the most biodiverse regions of the country and the world, located between 0°21' and 2°40' north latitude and 76°50' and 79°02' west longitude, with approximately 33,268 km<sup>2</sup> (Canchala et al., 2020; Ocampo-Marulanda et al., 2022). The Pacific region (14,754 km<sup>2</sup>) accounts for 52%, the Andean region (15,466 km<sup>2</sup>) for 40% and the Amazon region (3,048 km<sup>2</sup>) for the remaining 8% (See Figure 1) (Gobernación de Nariño, 2019). In addition, it has a privileged geostrategic position due to its proximity to the tropical Pacific Ocean, the Andes Mountains, and the Colombian-Ecuadorian border (Canchala et al., 2019).

Meteopo

logica

#### **2.2**) Data

In this research, time series of maximum (TMAX), minimum (TMIN), and mean temperature (TMED) in °C, humidity (RH) in %, sunshine hours (SBH) in hours, and height in meters above sea level (COTA) were considered as regressor variables on a monthly scale from 1983 to 2017. Data from 10 climatological stations across southwestern Colombia (see Figure 1) were provided by IDEAM (see Figure 1). The missing data in the time series were less than 25%. They were estimated using Non-Linear Principal Component Analysis (NLPCA), a methodology suggested by Scholz et al. (2005), and applied in hydroclimatology by Canchala et al. (2019).

Wind speed information was only available for three stations, Aeropuerto Antonio Nariño, El Encano, and Obonuco. The imputation of missing data was performed as mentioned in the Methods.

#### 2.3) Methods

Table 1 shows some methods for estimating ETo. FAO recommends using the *FAO-PM56* method to determine ETo without lysimeters. The estimation involves a wide range of variables, mean air temperature, relative humidity, solar radiation, and wind speed. Empirical models have been developed to estimate ETo with fewer climatic variables. For example, *Hargreaves* developed a model based on maximum, mean, minimum temperature, and solar radiation. *Turc* based his model on mean temperature, relative humidity, and net solar radiation. *Thornthwaite* proposed a model based on mean temperature and annual heat index, and Jaramillo based his model on the relationship between altitude and evapotranspiration.

The assessed performance metrics are presented in Table 2. The following metrics were used to evaluate the performance and precision of the alternative methods for estimating ETo, compared to *FAO-PM56*. RMSE, which characterizes the variance of the error (Rodrigues and Braga, 2021). MAE takes the absolute value of the difference between ETo values (Choi and Jeon, 2018). MBE measures the average error magnitude of the observed and estimated data (Goh et al., 2021). NSE is used to evaluate the predictive ability of hydrological models (Nash and Sutcliffe 1980; Knoben et al., 2019; Adnan et al., 2021). Finally, R measures the linear relationship between estimated and calculated values (Laqui et al., 2019).

The Machine learning methods used in this research were:

– Multiple Linear Regression (MLR). Regression analysis is a statistical technique belonging to the class of supervised statistical learning methods that allows investigating and modelling the relationship between a response variable and one or multiple predictor variables. An advantage of multiple linear regression is that it allows to evaluation of the effect of each predictor variable in the presence of the other variables (Montgomery et al., 2002). The multiple linear regression model is presented in Equation 1.

$$y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \ldots + \beta_k X_k + e$$
 (1)

Where

y = Response variable

 $\beta o =$  Intercept with the y-axis

 $\beta$ 1,...,Bk = partial regression coefficients that measure the expected change in the response variable for each unit change in the predictor variable X (j=1,2,...,k), when all other regressor variables are held constant.

e = Random component of error that must comply with the assumptions of normality, zero mean, constant variance (homoscedasticity) (Goldfend and Quandt, 1965), and independence (Fox, 2016).

- Autoregressive Integrated Moving Average (ARIMA). The ARIMA model is a statistical methodology that allows describing the future behavior of a time series as a linear function of past data and errors due to chance, in addition to considering the possible inclusion of a seasonal component (Box et al., 1974). The model that allows considering regular (non-seasonal) effects in the series can be expressed as in Equation 2.

 $(1 - \Phi_1 B - \Phi_2 B^2 - \dots - \Phi_p B^p)(1 - B)^d \text{ yt} = (1 - \theta_1 B - \theta_2 B^2 - \dots - \theta_q B^q)$ (2)

Where p and q indicate the order of the autoregressive (AR) and moving average (MA) components respectively, and d indicates the order of the integrated component (I) to extract the possible sources

of the non-stationarity present in the series under analysis (Stellwagen and Tashman, 2013). The AR and MA terms can be identified using the simple autocorrelation functions (ACF) and partial autocorrelation functions (PACF) of the time series data. ARIMA models were used to model the temporal correlation presented in the MLR models.

– Artificial Neuronal Network (ANN). AANs are computational bio-inspired models based on biological neurons, which can store and retrieve data, classify patterns, realize input patterns to output patterns, and similar group patterns. These follow two learning processes, supervised and unsupervised (Tabari and Talaee, 2013). Multilayer perceptron (MLP) is a type of feed-forward ANN mainly used for supervised learning (Haykin, 1994) and models complex nonlinear processes in water resources and hydrology problems. The MLP is a perceptron network with more than one intermediate layer and is usually represented with an output. The perceptron uses a matrix to model a neural network and is mainly used to discriminate an input x to a single output value F(x) in that matrix (See Equation 3).

$$F(\mathbf{x}) = \begin{cases} 1 \text{ if } w \cdot x \cdot u > 0 \\ 0 \text{ otherwise} \end{cases}$$
(3)

The function F(x) has a binary value and is used primarily for sorting, w is a vector with an associated weight, and u is a "threshold" used to offset the activation function. The sum of the inputs to the neuron must produce a value greater than u to change the neuron from state 0 to 1.

A successful development for an ANN depends on several parameters, e.g., hidden layers, neurons in the hidden layer, learning rate, and activation function, among others. However, there is no guideline on how to build an ANN or how many neurons should be placed in the hidden layer to estimate the output (Murat and Serhat, 2018). Therefore, through trial-and-error tests, the parameters were varied until the combination with the lowest error and the highest possible R<sup>2</sup> was found. In this research multiple architectures were built, combining activation functions, such as: identity, tanh, logistic and relu with the optimizers lbgfs, adam and sgd. Furthermore, given the complexity and the requirements one hidden layer with three neurons and a learning rate of 0.01 was enough. Through trial-and-error test, the best performance model with 1000 iterations was determined, and the best model was: tanh function activation, lbgfs optimizer, one hidden layer, three neurons and a learning rate of 0.01. The training algorithm of the ANN was 80% of the data and 20% to validate.

In selecting regressor variables for the ARIMA and MLR models, the VIF was used to avoid multicollinearity among the variables (Montgomery et al., 2002). In the case of ANNs, this technique performs nonlinear computational procedures which are not affected by multicollinearity because

## Meteoro logica

## Artículo en edición

they tend to be overparameterized, i.e., the same algorithm updates the weights associated with redundant variables to have no impact on the final solution (De Veaux and Ungar, 1994).

Linear Principal Component Analysis (PCA) was applied to formulate one of the ARIMA models. PCA is one of the most widely used statistical techniques to reduce dimensionality and preserve the most significant amount of information in a data set (Jolliffe, 2002), allowing the elimination of possible multicollinearity between regressor variables. The principal components between TMAX, TMED, and TMIN were estimated in this case.

Lee et al. (2012) mentioned that ETo is affected by topographic factors such as altitude because as altitude increases, there is a decrease in solar radiation and wind speed. Therefore, to represent the monthly ETo, an interpolation process was performed using the cokriging technique using altitude as an exogenous variable and with a spherical semivariogram. Basconcillo et al. (2017) and Cerón et al. (2021) suggest that with this technique, better correlations are obtained with monthly temperature and precipitation; considering that the former is one of the variables most correlated with ETo, it will be possible to spatialize ETo in southwestern Colombia.

The general methodology is presented in Figure 2. Initially, the information provided by IDEAM was compiled, then the exploratory data analysis was carried out, and the missing data were estimated. Subsequently, the ETo of the three stations was estimated using the *Hargreaves, Thornthwaite, Cenicafé*, and *Turc* methods. Then, the ETo methods were evaluated using five performance metrics, Nash-Sutcliffe model efficiency coefficient (NSE), Pearson correlation coefficient I, root mean square error (RMSE), mean absolute error (MAE) and mean bias error (MBE), to select the best-fit method concerning for *FAO-PM56*. Next, Spearman's correlation coefficient was applied to find the variables with the highest correlation with ETo and the variance inflation factor (VIF) to reduce multicollinearity among the regressor variables of the proposed statistical models. Finally, three models were built for each machine learning technique (Artificial Neural Networks – ANN, Multiple Linear Regression – RLM, and Autoregressive Integrated Moving Average Model – ARIMA), i.e., nine models, which were evaluated by five performance metrics (NSE, R, RMSE, MAE and MBE) to estimate *FAO-PM56* with scarce information.

Then, multi-year monthly averages of all variables were estimated, and the annual value of precipitation was reported. Subsequently, the Jarque-Bera normality test was performed, which tests whether a data set presents the skewness and kurtosis of a normal distribution (Jarque and Bera, 1982). The test was used to determine parametric or non-parametric statistical tests to calculate the correlation between climatic variables. Empirical models have been developed to estimate Eto with

fewer climatic variables.

#### 3) RESULTS AND DISCUSSION

The descriptive statistics of the climatic variables were estimated and presented in Table 3. It can be observed that the altitude of the stations is in the mountainous zone, except for Barbacoas, which is in the Pacific plain, and Monopamba, located in the Amazon jungle (see Figure 1). The precipitation values ranged between 888 and 6,927 mm/year, consistent results considering that the department includes the Andean, Amazon, and Pacific regions, the latter being one of the rainiest regions in the world. The TMAX ranges between 32.5 °C and 18.8 °C, the TMED between 11.7 °C and 26.2 °C, and the TMIN between 4.8 °C and 20 °C. The RH presented values above 75%, with 89.5% being the highest reported at the Barbacoas station. Finally, the SBH reported values between 2.1 and 5.4 hours. The indicators shown in Table 3 were calculated with the complete time series, with a previous estimation of missing data using NLPCA.

The reconstruction errors were, 7 mm month<sup>-1</sup> for precipitation, 0.16 °C month<sup>-1</sup> for maximum temperature, 0.07 °C month<sup>-1</sup> for mean temperature, 0.16 °C month<sup>-1</sup> for minimum temperature, 0.55 % month<sup>-1</sup> for relative humidity and 0.10 hours month<sup>-1</sup> for sunshine hours. These results show very low error variance in magnitude, demonstrating that the imputation method accurately estimated the missing data in the time series.

Figure 3 shows a graphical comparison of ETo methods with *FAO-PM56* for the time series of the Aeropuerto Antonio Nariño, El Encano and Obonuco stations to find the method with the most well-adjustment to *FAO-PM56*. *Hargreaves* method was the only one that overestimated ETo, since the others tended to underestimate it. The *Turc* method was similar in shape and magnitude to the *FAO-PM56* method at El Encano station, while at Obonuco, the most well-adjusted method was *Hargreaves*. At the Aeropuerto Antonio Nariño station, it was observed that both methods were adjusted in shape and magnitude.

In this sense, the performance metrics were estimated to validate the most adjusted method concerning *FAO-PM56*, and the results are presented in Table 4. The average RMSE performance metrics ranged from 16.9 to 50.4 mm month<sup>-1</sup>, MAE from 14.2 to 48.7 mm month<sup>-1</sup>, MBE from 10.3 to 48.7 mm month<sup>-1</sup>, R from 0.80 to 0.90 and NSE from 0.64 to 0.81. The results of the *Turc* method show that it was the most well-adjusted concerning *FAO-PM56* by the reported metrics (MAE= 14.2 mm month<sup>-1</sup>, MBE=10.3 mm month<sup>-1</sup> and RMSE= 16.9 mm month<sup>-1</sup>). However, the R and NSE metrics (0.90 and 0.81) show *Hargreaves* as the best. Trajkovic and Kolakovic (2009a) state that *Turc* 

## Meteoro logica

## Artículo en edición

method overestimates ETo at windless locations and underestimates at windy locations; for example, Figure 3 shows underestimation in Aeropuerto Antonio Nariño, Obonuco and El Encano stations located at heights above 1700 m. These results agree with Poveda et al. (2007), who regionalized evapotranspiration in Colombia and observed that the *Turc* and *Morton* methods are the most appropriate; and Fisher & Pringle (2013), who used three alternative methods in a humid region in the United States (Mississippi) and concluded that Turc consistently underestimates ETo.

Other research reports that the *Turc* method has historically performed well under humid (Jensen et al., 1990; Trajkovic and Kolakovic, 2009b; Fisher & Pringle, 2013; Birara et al., 2020) and tropic conditions (Tukimat et al., 2012; Lima et al., 2019; Santos et al., 2019; Monteiro et al., 2021). One of the possible reasons is that the method was initially developed under wet conditions (southern France) (Tabari, 2010; Ahmadi and Javanbakht 2020; Diouf et al., 2016). In a semiarid region in Senegal Diouf et al. (2016) concluded that the *Turc* method showed similar high accuracy ( $R^2 > 0.80$ ) reported in this research. Similar findings were observed in Malaysia, where Goh et al. (2021) concluded that the Turc method provides the closest results to FAO-PM56 in subhumid and humid climate conditions in the absence of data, as the  $R^2$  and the MBE reported one of the highest  $R^2$  (0.81) and the MBE results were closer to zero for the monthly ETo estimation concerning FAO-PM56. The Same happened in Brazil, where Santos et al. (2019) concluded that the best option when meteorological data is unavailable is the Turc method given the R results (0.90) concerning FAO-PM56. In contrast, Monteiro et al. (2021) suggest that the ETo estimation should give priority to Turc Method regardless of the season and climatic conditions. From these results, it can be concluded that in the absence of information to estimate FAO-PM56, the Turc method is the best alternative to estimate ETo in southwestern Colombia.

When the normality test (Jarque-Bera) was applied to the regressor variables, a p-value of 0.00 was obtained, which means that the data did not follow a normal distribution. Therefore, the nonparametric Spearman correlation coefficient was applied between the climatic variables and *FAO-PM56*. The results were, COTA -0.84, RH -0.69, SBH 0.87, TMAX 0.88, TMED 0.84 and TMIN 0.66, which suggest the highest positive correlation concerning *FAO-PM56* are TMAX, SBH and TMED, and the highest negative correlation was COTA. However, SBH is a complex variable to obtain due to its high costs and technical complexities (Laidi, 2018). TMED results in a lower saturation pressure, hence a lower vapour pressure deficit, which results in a lower estimate of ETo (Allen et al., 1998). Hence, TMAX and COTA were considered the most crucial regressor variables for constructing the ARIMA and MLR models, as shown in Table 5, where the VIF results are also

presented to identify any possible multicollinearity.

The TMIN and RH presented the lowest correlation coefficients (0.66 and -0.69), which indicates that they are not the best choice as regressor variables. Multicollinearity was evidenced for TMAX, TMED and COTA due to the high values (> 5) in the VIF. These results are congruent, considering that both temperatures are correlated and have an inversely proportional relationship with altitude. Therefore, the TMAX variable was prioritized for the construction of the models since it was the one that reported the highest correlation coefficient (0.88).

Figure 4 presents the graphical comparison between the best machine learning models, MLR-M2, ANN-M1 and ARIMA-M3, during 1983-2017, reported in Table 6. The ARIMA-M3 method was the closest in magnitude and shape for the Aeropuerto Antonio Nariño station. The MLR-M2 model was the one that reported the highest accuracy in the extreme values for the three stations, evidencing it in the maximum and minimum peaks. However, these methods cannot be extrapolated to other climatological stations because they were calibrated with specific COTA and TMAX conditions, and stations. The ANN-M1 model underestimates ETo at Aeropuerto Antonio Nariño, although it reported higher accuracy at El Encano and Obonuco stations. However, it is necessary to perform a quantitative evaluation to validate the graphical results. Therefore, Table 6 presents the performance metrics results for the nine proposed machine learning models.

The performance metrics showed that the average of 9 models ranged in RMSE from 4.1 to 8.2 mm month<sup>-1</sup>, MAE from 3.2 to 6.6 mm month<sup>-1</sup>, MBE from -0.1 to 0.3 mm month<sup>-1</sup>, NSE from 0.84 to 0.96 and r from 0.92 to 0.98. Of the 9 models constructed, the ARIMA-M3 model reported the best results in terms of error (RMSE=4.1 mm month<sup>-1</sup>, MAE=3.2 mm month<sup>-1</sup>, MBE= -0.1 mm month<sup>-1</sup>, NSE=0.96 and R=0.98). This result suggests that using this model would allow estimating ETo more accurately in error and correlation. However, these can only be used if there is prior information on the response variable (*FAO-PM56*) due to its autoregressive component, making it difficult to use in places with scarce information. Although the ANNs did not report the best metrics, they can be used as an alternative for estimating *FAO-PM56* in places with scarce information since it is not necessary to know the response variable for its estimation.

These results agree with what has been reported in other research (Zanetti et al., 2008; Alves et al., 2017; Nema et al., 2017; Laqui et al., 2019; Granata et al., 2020; Meneses et al., 2020), where it is shown that the application of ANNs using MLP with few regressor variables allows estimating ETo more accurately. Moreover, Feng et al. (2018) and Shiri (2017) suggest that machine learning models outperformed empirical equations, which is evident in the results of the performance metrics, as the

# Meteoro logica

## Artículo en edición

*Turc* method goes from having RMSE from 16.9 to 7.3 mm month<sup>-1</sup>, MAE from 14.2 to 5.9 mm month<sup>-1</sup>, MBE from 10.3 to 0.1 mm month<sup>-1</sup>, R from 0.81 to 0.87 and NSE from 0.65 to 0.94. It is crucial select TMAX, SBH, and COTA to achieve efficient ANN models in regions with scarce information. To improve the error and efficiency of the ANN, we recommend adding wind speed, given that Maček et al. (2018) suggest that it accounts for a significant contribution to the aerodynamic component. However, unfortunately, this is the most complex variable to get data due to the lack of stations in the country.

Even though there are no previous monthly scale studies on this methodology in Colombia, Laqui et al. (2019) investigated in a similar context (Peruvian highlands), i.e., climatological stations with scarce information located at high altitudes (3819 and 4660 masl). Although the scale was a daily scale, the results showed that ANNs allow estimating with reasonable accuracy at high altitude stations with scarce information. Pinos et al. (2020) assesses 30 models for the estimation of daily ETo in two weather stations with limited data in a wet Andean paramo ecosystem (southern Ecuador). Their results suggest that the ANNs outperformed the empirical models and accurately estimated ETo in super-humid conditions. It is agreed with different studies developed in humid (Ayaz et al., 2021), subhumid (Nema et al., 2017), arid (Tabari and Talaee, 2013), semiarid (Ayaz et al., 2021) and wetland areas (Granata et al., 2020) around the world where ANN were used to estimate ETo with few input variables.

In northern Greece, Antonopoulos & Antonopoulos (2017) build 3 ANNs models and concluded that any model that uses temperature and radiation as inputs should be able to estimate the ETo sufficiently. Mohawesh (2013) proved several ANNs in 3 stations across the Jordan valley and concluded that the overall results suggest that temperature based ANNs can be used when there is insufficient data. In Serbia, Petković et al. (2015) developed an adaptive neuro-fuzzy inference system and concluded that the maximum relative humidity and maximum air temperature are the most influential optimal. The same happened in Brazil, where Ferreira (2019) concluded that the relative humidity and temperature increased the capacity of the ANNs to estimate ETo adequately. Thereby, the results shown in this research represent an option to substitute the *FAO-PM56* method in regions with scarce information; since the amount of information required by this method could be a limiting factor for its application.

Figure 5 shows the results of the ETo estimation with the ANN-M1 model for the seven stations for which *FAO-PM56* could not be estimated. It shows that most results present the same shape and magnitude except for Barbacoas, which is congruent considering that it is in the Pacific region with

marked differences concerning the Andean region, where most other stations are located. Interpolation with ordinary cokriging with the spherical semivariogram from the stations of this research to know the annual ETo in southwestern Colombia was performed. The results in Figure 6 show that the highest values are found in the Pacific region. Lower values are observed as one approaches the Andean region and reaches the Amazon region. A noteworthy result is that the reported values of ETo in the Andean region in comparison with precipitation could indicate a water deficit.

#### 4) CONCLUSIONS

The results show that this proposed machine learning models allow a precise estimation of ETo in southwestern Colombia with scarce information since the performance metrics were better than those reported by the best-fitted empirical method (*Turc*). However, as these were far from the ideal value, it was decided to build machine learning techniques to reduce the error associated with ETo estimation. Initially, four regressor variables were considered for estimation, and given the high correlation with *FAO-PM56* (Spearman correlation coefficient) with TMAX (0.88) and COTA (-0.84) and easiness of obtaining these variables, the machine learning model was built prioritizing TMAX and COTA over the others. More input variables give accurate information about ETo estimation, as many researchers suggest (Rivas and Caselles, 2004; Nema et al., 2017; Choi and Jeon, 2018); yet one of the purposes of this research is to develop an alternative model with few inputs variables to ease application to adequately estimate ETo in the absence of data as in southwestern Colombia.

Hence, ETo was calculated with nine machine learning models, and it was determined that the ARIMA-M3 and MLR-M2 models presented the best performance, because it was calibrated with specific conditions like COTA, TMAX (exogenous variables), and stations. However, there are restrictions on their use since they cannot be extrapolated outside the study area. Therefore, the ANN-M1 model was used as an alternative method to estimate ETo in southwestern Colombia with scarce information, considering previous successful studies, their unrestricted application, and good performance metrics, and that does not require knowing the response variable for its estimation since it works as a black-box model.

Metrics performance of the ANN-M1 model concerning those calculated by the *Turc* method is better, as it goes from having an RMSE of 16.9 to 7.3 mm month<sup>-1</sup>, an MAE of 14.2 to 5.9 mm month<sup>-1</sup>, an MBE of 10.3 to 0.1 mm month<sup>-1</sup>, an R of 0.81 to 0.87 and an NSE of 0.65 to 0.94. These results suggest that using the ANN-M1 model allows a more accurate estimation of ETo in places with little

information at high altitudes, which allows considering it as a methodology to be used in forecasts or to improve the understanding of future hydroclimatic events to reduce the uncertainty generated.

The ANNs tend to underestimate the ETo results slightly; this may be because the developed network did not adequately assimilate the spatial variation with the current information. Considering that the results obtained are reasonable, it is suggested to improve the data source by installing new climatological stations to better train the ANNs. It is also necessary to better understand the spatial-temporal variation of the climatic variables and, thus, to have better planning and management in managing the water resources of the southwestern part of Colombia.

The results presented here contributed to validating the idea of the application of machine learning techniques to estimate ETo in places with scarce information. This fact provides, easy and accurate information to agriculturists and stakeholders to develop programs that provide and enhance water resources management to achieve the proposed sustainable development goals and ensure food security and sovereignty. Future studies could emphasize developing machine learning models (ANN) to estimate and forecast the ETo in different climate conditions, e.g., arid and humid, on a monthly scale.

#### Acknowledgment

The authors thank the Universidad del Valle for financing the research project CI 21010 and the Colombia Science Ministry for funding the research project "Análisis de eventos extremos de precipitación asociados a variabilidad y cambio climático para la implementación de estrategias de adaptación en sistemas productivos agrícolas de Nariño". Furthermore, the authors thank the IREHISA, GUIA, and INFERIR research groups at Universidad del Valle for developing this research paper. Finally, the authors express their gratitude to IDEAM for providing the monthly information of climatological variables in southwestern Colombia. The last author has received research support from Fondo Nacional de Financiamiento para la Ciencia, la Tecnología y la Innovación Francisco José de Caldas – MINCIENCIAS through the "Convocatoria No 891 de 2020 para el fortalecimiento de vocaciones y formación en CteI para la reactivación económica en el marco de la postpandemia 2020". The authors thank the two anonymous reviewers for their helpful comments.

#### 5) **REFERENCES**

Adnan, R. M., Heddam, S., Yaseen, Z. M., Shahid, S., Kisi, O., Li, B., 2021: Prediction of potential evapotranspiration using temperature-based heuristic approaches. Sustainability 13(1):1–21.

https://doi.org/10.3390/su13010297

Ahmadi, S. H., Javanbakht, Z., 2020: Assessing the physical and empirical reference evapotranspiration (ETo) models and time series analyses of the influencing weather variables on ETo in a semiarid area. Journal of Environmental Management 276:111278. https://doi.org/10.1016/j.jenvman.2020.111278

Allen, R. G., Pereira, L. S., Raes, D., Smith, M., 1998: Crop evapotranspiration – guidelines for computing crop water requirements – FAO Irrigation and drainage 56, pp. 1-322.

Alves, W. B., Rolim, G. D., Aparecido, L. E., 2017: Reference evapotranspiration forecasting by artificial neural networks. Engenharia Agricola 37(6):1116–1125. https://doi.org/10.1590/1809-4430-eng.agric.v37n6p1116-1125/2017

Antonopoulos, V. Z., Antonopoulos, A. V., 2017: Daily reference evapotranspiration estimates by artificial neural networks technique and empirical equations using limited input climate variables. Computers and Electronics in Agriculture 132, 86–96. https://doi.org/10.1016/j.compag.2016.11.011

Ayaz, A., Rajesh, M., Singh, S. K., Rehana, S., 2021: Estimation of reference evapotranspiration using machine learning models with limited data. AIMS Geosciences 7(3):268–290. https://doi.org/10.3934/geosci.2021016

Barco, J., Cuartas, A., Mesa, O., Poveda, G., Vélez, J. I., Hoyos, C., Mejía, J. F., Botero, B., 2000: Estimación de la evapotranspiración en Colombia. Avances en recursos hidráulicos, 7:43–51.

Basconcillo, J., Duran, G. A., Fransisco, A., Abastillas, R., Hilario, F., Juanillo, E., Solis, A. L., Lucero, A. J., Maratas, S.-L., 2017: Evaluation of Spatial Interpolation Techniques for Operational Climate Monitoring in the Philippines. SOLA 13:114–119. https://doi.org/10.2151/sola.2017-021

Biggs, T. W., Marshall, M., Messina, A., 2016: Mapping daily and seasonal evapotranspiration from irrigated crops using global climate grids and satellite imagery: Automation and methods comparison. Water Resour. Res. 52:7311–7326. https://doi.org/10.1002/2016WR019107

Birara, H., Mishra, S. K. & Pandey, R. P, 2020: Comparison of methods for evapotranspiration computation in the Tana Basin, Ethiopia, Volume 97, In: Hydrological Extremes. Springer, pp. 1-472. Blaney H., Criddle, W., 1950: Determining water requirements in irrigated areas from climatological and

irrigation data.

Bouznad, I. E., Guastaldi, E., Zirulia, A., Brancale, M., Barbagli, A., Bengusmia, D., 2020: Trend analysis and spatiotemporal prediction of precipitation, temperature, and evapotranspiration values using the ARIMA models: case of the Algerian Highlands. Arabian Journal of Geosciences 13(24). https://doi.org/10.1007/s12517-020-06330-6

Box, G. E. P., Jenkins, G. M., Reinsel, G. C., Ljung, G. M., 1994: Time series forecasting: forecasting and control. Fifth edition. Prentice Hall. Englewood Cliff. New Jersey, pp. 1-422.

Canchala, T., Alfonso-Morales, W., Carvajal-Escobar, Y., Cerón, W. L., Caicedo-Bravo, E., 2020: Monthly Rainfall Anomalies Forecasting for Southwestern Colombia Using Artificial Neural. Water 12(9):2628. https://doi.org/10.3390/w12092628

Canchala, T., Ocampo-Marulanda, C., Alfonso-Morales, W., Carvajal-Escobar, Y., Ceron, W. L., Caicedo-Bravo, E. (2022). Techniques for monthly rainfall regionalization in southwestern Colombia. Anais da Academia Brasileira de Ciências, 94. <u>https://doi.org/10.1590/0001-3765202220201000</u>

Canchala-Nastar, T., Carvajal-Escobar, Y., Alfonso-Morales, W., Loaiza, W. C., Caicedo, E., 2019: Estimation of missing data of monthly rainfall in southwestern Colombia using artificial neural networks. Data in Brief 26. https://doi.org/10.1016/j.dib.2019.104517

Cannarozzo, M., Noto, L. V., Viola, F. Spatial distribution of rainfall trends in Sicily (1921-2000)., 2006: Physics and Chemistry of the Earth 31(18):1201–1211. https://doi.org/10.1016/j.pce.2006.03.02

Castañeda, L., Rao, P., 2005: Comparison of methods for estimating reference evapotranspiration in Southern California. Journal of Environmental Hydrology 1–10. http://hydroweb.com/jeh/jeh2005/castaneda.pdf

Cerón, W., Andreoli, R., Kayano, M., Canchala, T., Carvajal-Escobar, Y., Souza, R., 2021: Comparison of spatial interpolation methods for annual and seasonal rainfall in two hotspots of biodiversity in South America. Annals of the Brazilian Academy of Sciences 93(1). https://doi.org/10.1590/0001-3765202120190674

Choi, Y., Jeon, J., 2018: Comparison of Artificial Neural Network and Empirical Models to Determine

Daily Reference Evapotranspiration. Journal of the Korean Society of Agricultural Engineers 60(6):43–54. https://doi.org/10.5389/KSAE.2018.60.6.043

Cobaner, M., 2010: Evapotranspiration estimation by two different neuro-fuzzy inference systems. Journal of Hydrology 398(3–4):292–302, https://doi.org/10.1016/j.jhydrol.2010.12.030

De Veaux R. D., Ungar, L. H., 1994: Multicollinearity: A tale of two nonparametric regressions. Lecture Notes in Statistics 393-402. https://doi.org/10.1007/978-1-4612-2660-4\_40

Diouf, O. C., Weihermüller, L., Ba, K., Faye, S. C., Faye, S., Vereecken, H., 2016: Estimation of Turc reference evapotranspiration with limited data against the Penman-Monteith Formula in Senegal. Journal of Agriculture and Environment for International Development 110(1), 117–137. https://doi.org/10.12895/jaeid.20161.417

Feng Y., Peng Y., Cui N., 2017: Modeling reference evapotranspiration using extreme learning machine and generalized regression neural network only with temperature data. Comput Electron Agric 136, 71–78. https://doi.org/10.1016/j.compag.2017.01.027

Ferreira, L. B., da Cunha, F. F., de Oliveira, R. A., Fernandes Filho, E. I., 2019: Estimation of reference evapotranspiration in Brazil with limited meteorological data using ANN and SVM – A new approach. Journal of Hydrology 572, 556–570. https://doi.org/10.1016/j.jhydrol.2019.03.028

Ferreira, L. B., França da Cunha, F., Zanetti, S. S., 2021: Selecting models for the estimation of reference evapotranspiration for irrigation scheduling purposes. PloS ONE 16. https://doi.org/10.1371/journal.pone.0245270

Fisher, D., Pringle, H., 2013: Evaluation of alternative methods for estimating reference evapotranspiration. Agricultural Sciences 04(08):51–60. https://doi.org/10.4236/as.2013.48a008

Fonseca-Luengo, D., Lillo-Saavedra, M., Lagos, L. O., García-Pedrero, A., Gonzalo-Martín, C., 2018: Use of machine learning to improve the robustness of spatial estimation of evapotranspiration. Lecture Notes in Computer Science 10657:237–245. https://doi.org/10.1007/978-3-319-75193-1\_29

Fox, J., 2016: Applied Regression Analysis and Generalized Linear Models. 3<sup>rd</sup> edition, SAGE Publications, pp. 1-817.

Gautam, R., Sinha, A. K., 2016: Time series analysis of reference crop evapotranspiration for Bokaro District, Jharkhand, India. Journal of Water and Land Development 30(1):51–56. https://doi.org/10.1515/jwld-2016-0021

Gobernación de Nariño, 2019: Plan participativo de desarrollo departamental 2016-2019. https://publicadministration.un.org/unpsa/Portals/0/UNPSA\_Submitted\_Docs/Plan%20de%20Desarrollo %20Departamental%20Nari%C3%B1o%20Coraz%C3%B3n%20del%20Mundo..pdf?ver=2018-11-29-171310-447, accessed on 01/05/2022.

Goh, E. H., Ng, J. L., Huang, Y. F., Yong, S. L. S., 2021: Performance of potential evapotranspiration models in Peninsular Malaysia. Journal of Water and Climate Change 00(0). https://doi.org/10.2166/wcc.2021.018

Goldfeld, S., Quandt, R., 1965: Some Tests for Homoscedasticity. Journal of the American Statistical Association 60:539–547. https://doi.org/10.1080/01621459.1965.10480811

Granata, F., Gargano, R., de Marinis, G., 2020: Artificial intelligence based approaches to evaluate actual evapotranspiration in wetlands. Science of the Total Environment 703:135653. https://doi.org/10.1016/j.scitotenv.2019.135653

Hamdi, M. R., Bdour, A. N., Tarawneh, Z. S., 2008: Developing Reference Crop Evapotranspiration Time Series Simulation Model Using Class a Pan : A Case Study for the Jordan. Jordan Journal of Earth and Environmental Sciences 1(1):33–44. https://eis.hu.edu.jo/deanshipfiles/pub10431145.pdf

Hargreaves, H. G., Samani, Z. A. Reference crop evapotranspiration from temperature., 1985: Applied Eng. In Agric 1(3).

Haykin, S., 1994: Neural Networks: A Comprehensive Foundation, Volume: 3, Prentice Hall PTR, pp. 1-823.

https://cdn.preterhuman.net/texts/science\_and\_technology/artificial\_intelligence/Neural%20Networks%2 0-%20A%20Comprehensive%20Foundation%20-%20Simon%20Haykin.pdf

Huizhi, L., Jianwu, F., 2012: Seasonal and interannual variations of evapotranspiration and energy exchange over different land surfaces in a semiarid area of China. Journal of Applied Meteorology and

Climatology 51(10): 1875-1888. https://doi.org/10.1175/JAMC-D-11-0229.1

Huo, Z., Feng, S., Kang, S., Dai, X., 2012: Artificial neural network models for reference evapotranspiration in an arid area of northwest China. Journal of Arid Environments 82:81–90. https://doi.org/10.1016/j.jaridenv.2012.01.016

Igbadun, H. E., Mahoo, H. F., Tarimo, A. K. P. R., Salim, B. A., 2006: Crop water productivity of an irrigated maize crop in Mkoji sub-catchment of the Great Ruaha River Basin, Tanzania. Agricultural Water Management 85(1–2):141–150. https://doi.org/10.1016/j.agwat.2006.04.003

Jaramillo, A. Evapotranspiración de referencia en la Región Andina de Colombia. Cenicafé, 2006:57(4):288–298. https://www.cenicafe.org/es/publications/arc057(04)288-298.pdf

Jarque, C., Bera, A., 1980: Efficient tests for normality, homoscedasticity, and serial independence of regression residuals. Economics Letters 6:255–259. http://l.academicdirect.org/Horticulture/GAs/Refs/Jarque&Bera 1980.pdf

Jensen, M. E., Burman R. D., Allen, R. G., 1990: Evapotranspiration and irrigation water requirements. ASCE Manual and Report No. 70, pp. 1-744.

Jing, W., Yaseen, Z. M., Shahid, S., Saggi, M. K., Tao, H., Kisi, O., Salih, S. Q., Al-Ansari, N., Chau, K. W.,2019: Implementation of evolutionary computing models for reference evapotranspiration modeling: short review, assessment and possible future research directions. Engineering Applications of Computational Fluid Mechanics 13(1):811–823. https://doi.org/10.1080/19942060.2019.1645045

Jolliffe, I., 2002: Principal Component Analysis. Springer, pp. 1-518.

Knoben, W. J. M., Freer, J. E., Woods, R. A., 2019: Technical note: Inherent benchmark or not? Comparing Nash-Sutcliffe and Kling-Gupta efficiency scores. Hydrology and Earth System Sciences 23(10)4323–4331. https://doi.org/10.5194/hess-23-4323-2019

Kumar, M., Raghuwanshi, N. S., Singh, R., Wallender, W. W., Pruitt, W. O., 2002: Estimating of Evapotranspiration Using Artificial Neural Network. Journal of Irrigation and Drainage Engineering 128:224–233. https://doi.org/10.1061/(ASCE)0733-9437(2002)128:4~224!4

Łabędzki, L., Bąk, B., Smarzyńska, K., 2014: Spatio-temporal variability and trends of Penman-Monteith reference evapotranspiration (FAO-56) in 1971-2010 under climatic conditions of Poland. Polish Journal of Environmental Studies 23(6):2083–2091. https://doi.org/10.15244/pjoes/27816

Laidi, M., Hanini, S., El Hadj, A., 2018: Novel approach for estimating monthly sunshine duration using artificial neural networks: A case study. Journal of Sustainable Development of Energy, Water and Environment Systems 6(3):405–414. https://doi.org/10.13044/j.sdewes.d6.0226

Landeras, G., Ortiz-Barredo, A., López, J. J., 2008: Comparison of artificial neural network models and empirical and semi-empirical equations for daily reference evapotranspiration estimation in the Basque Country (Northern Spain). Agricultural Water Management 95(5):553–565. https://doi.org/10.1016/j.agwat.2007.12.011

Laqui, W., Zubieta, R., Rau, P., Mejía, A., Lavado, W., Ingol, E., 2019: Can artificial neural networks estimate potential evapotranspiration in Peruvian highlands?. Modeling Earth Systems and Environment 5(4):1911–1924. https://doi.org/10.1007/s40808-019-00647-2

Lee, B.-Y., Yang, S.-K., Kwon, K.-H., Kim, J.-B., 2012: The Effect of Evapotranspiration by Altitude and Observation of Lysimeter. Journal of Environmental Science International 21(6):749–755. https://doi.org/10.5322/jes.2012.21.6.749

Lima, J. G., Viana, P. C., Sobrinho, J. E., Couto, J. P., 2019: Comparison of ETo estimation methods and sensitivity analysis for different Brazilian climates. Irriga 2019:24(3), 538–551. https://doi.org/10.15809/irriga.2019v24n3p538-551

Liu, Q., Yang, Z., Cui, B., 2008: Spatial and temporal variability of annual precipitation during 1961-2006 in Yellow River Basin, China. Journal of Hydrology 361(3–4):330–338. https://doi.org/10.1016/j.jhydrol.2008.08.002

Maček, U., Bezak, N., Šraj, M., 2018: Reference evapotranspiration changes in Slovenia, Europe.AgriculturalandForestMeteorology260–261:183–192.https://doi.org/https://doi.org/10.1016/j.agrformet.2018.06.014

Mendoza C., C. J., Peña Q., A. J., 2021: Reference evapotranspiration estimation by different methods for the sucroenergy sector of Colombia. Revista Brasileira de Engenharia Agricola e Ambiental 25(9):583–

590. https://doi.org/10.1590/1807-1929/agriambi.v25n9p583-590

Meneses, K. C., Aparecido, L. E., Meneses, K. C., Farias M. F., 2020: Estimating potential evapotranspiration in maranhão state using artificial neural networks. Revista Brasileira de Meteorologia 2020:35(4):675–682. https://doi.org/10.1590/0102-77863540072

Mohawesh, O. E., 2013: Artificial neural network for estimating monthly reference evapotransiration under arid and semi arid environments. Archives of Agronomy and Soil Science 59(1), 105–117. https://doi.org/10.1080/03650340.2011.603126

Moncayo, C., 2015: Valle del Cauca, Antioquia y Nariño, departamentos líderes en producción., https://incp.org.co/valle-del-cauca-antioquia-y-narino-departamentos-lideres-enproduccion/#:~:text=Nari%C3%B10%20tiene%20una%20producci%C3%B3n%20significativa,banano% 20(12%2C8%25, accessed on 31.03.2022.

Monteiro, A. F., Martins, F. B., Torres, R. R., de Almeida, V. H., Abreu, M. C., Mattos, E. V., 2021: Intercomparison and uncertainty assessment of methods for estimating evapotranspiration using a high-resolution gridded weather dataset over Brazil. Theoretical and Applied Climatology 146(1–2), 583–597. https://doi.org/10.1007/s00704-021-03747-1

Montgomery, D., Peck E., Vining, G., 2002: Introducción al análisis de regresión lineal. Compañía Editorial Continental. México D.F, pp- 1-590.

Mossad, A., Alazba, A. A., 2016: Simulation of temporal variation for reference evapotranspiration under arid climate. Arabian Journal of Geosciences 9(5). https://doi.org/10.1007/s12517-016-2482-y

Murat, A., Serhat, Ö., 2018: Artificial intelligence (AI) Studies in Water Resources. Natural and Engineering Sciences 3(2), 187–195. https://www.researchgate.net/publication/325212839\_Artificial\_Intelligence\_AI\_Studies\_in\_Water\_Reso urces/fulltext/5afe2e41aca272b5d84a9e57/Artificial-Intelligence-AI-Studies-in-Water-Resources.pdf

Nash, J. E., Sutcliffe, J. V., 1980: River flow forecasting through conceptual models. Part I—A discussion of principles. Journal of Hydrology 10(3):282–290. https://doi.org/10.1016/0022-1694(70)90255-6

Nema, M. K., Khare, D., Chandniha, S. K., 2017: Application of artificial intelligence to estimate the



reference evapotranspiration in sub-humid Doon valley. Applied Water Science 7(7):3903–3910. https://doi.org/10.1007/s13201-017-0543-3

Ocampo-Marulanda, C., Fernández-Álvarez, C., Cerón, W. L., Canchala, T., Carvajal-Escobar Y., Alfonso-Morales W., 2022: A spatiotemporal assessment of the high-resolution CHIRPS rainfall dataset in southwestern Colombia using combined principal component analysis. Ain Shams Eng. J. 13(5):101739. https://doi.org/10.1016/j.asej.2022.101739

Petković, D., Gocic, M., Trajkovic, S., Shamshirband, S., Motamedi, S., Hashim, R., Bonakdari, H., 2015: Determination of the most influential weather parameters on reference evapotranspiration by adaptive neuro-fuzzy methodology. Computers and Electronics in Agriculture 114, 277–284. https://doi.org/10.1016/j.compag.2015.04.012

Pinos, J., Chacón, G., Feyen, J., 2020: Comparative Analysis of Reference Evapotranspiration Models with Application to The Wet Andean Páramo Ecosystem in Southern Ecuador. Meteorologica, 45(1):25–45. http://www.meteorologica.org.ar/wp-content/uploads/2020/03/v45n1a02-1.pdf

Poveda, G., Vélez, J. I., Mesa, O. J., Cuartas, A., Barco, J., Mantilla, R. I., Mejía, J. F., Hoyos, C. D., Ramírez, J. M., Ceballos, L. I., Zuluaga, M. D., Arias, P. A., Botero, B. A., Montoya, M. I., Giraldo, J. D., Quevedo, D. I., 2017: Linking Long-Term Water Balances and Statistical Scaling to Estimate River Flows along the Drainage Network of Colombia. Journal of Hydrologic Engineering 12(1):4–13. https://doi.org/10.1061/(asce)1084-0699(2007)12:1(4)

Priestley C. H. B. Taylor R. J., 1972: On the assessment of surface heat flux and evaporation using large-scale parameters. Mon. Weather Rev 100:2:81–92.

Quej, V. H., Almorox, J., Arnaldo, J. A., Moratiel, R., 2019: Evaluation of Temperature-Based Methods for the Estimation of Reference Evapotranspiration in the Yucatán Peninsula, Mexico. Journal of Hydrologic Engineering, 24(2):05018029. https://doi.org/10.1061/(asce)he.1943-5584.0001747

Ramírez, V. H., Mejía, A., Marín, E. V., Arango, R., 2011: Evaluation of models for estimating the reference evapotranspiration in Colombian Coffee Zone. Agronomía Colombiana, 29(1):107–114. https://repositorio.unal.edu.co/bitstream/handle/unal/39836/28641-102621-2-PB.pdf?sequence=1&isAllowed=y

Rivas, R., & Caselles, V., 2004: A simplified equation to estimate spatial reference evaporation from remote sensing-based surface temperature and local meteorological data. Remote Sensing of Environment 93(1–2):68–76. https://doi.org/10.1016/j.rse.2004.06.021

Rodrigues, G. C., Braga, R. P., 2021: Estimation of reference evapotranspiration during the irrigation season using nine temperature-based methods in a hot-summer mediterranean climate. Agriculture 11(2):1–15. https://doi.org/10.3390/agriculture11020124

Santos, L. da C., Cruz, G. H., Capuchinho, F. F., José, J. V., & dos Reis, E. F., 2019: Assessment of empirical methods for estimation of reference evapotranspiration in the Brazilian Savannah. Australian Journal of Crop Science 13(7), 1094–1104. https://doi.org/10.21475/ajcs.19.13.07.p1569E

Scholz, M., Kaplan, F., Guy, C. L., Kopka J. Selbig, J., 2005: Non-linear PCA: a missing data approach. Bioinformatics 21:3887-3895. https://doi.org/10.1093/bioinformatics/bti634

Shiri J., 2017: Evaluation of FAO56-PM, empirical, semi-empirical and gene expression programming approaches for estimating daily reference evapotranspiration in hyper-arid regions of Iran. Agric Water Manag 188, 101–114. https://doi.org/10.1016/j.agwat.2017.04.009

Stellwagen, E., Tashman, L., 2013: ARIMA : The Models of Box and Jenkins. Foresight: The International Journal of Applied Forecasting, 30:28–34.

Tabari, H., Talaee, P. H., 2013: Multilayer perceptron for reference evapotranspiration estimation in a semiarid region. Neural Computing and Applications 23(2):341–348. https://doi.org/10.1007/s00521-012-0904-7

Tabari, H., Talaee, P. H., 2013: Multilayer perceptron for reference evapotranspiration estimation in a semiarid region. Neural Computing and Applications 23(2):341–348. https://doi.org/10.1007/s00521-012-0904-7

Thornthwaite, C., Wilm, H., 1948: Report of the Committee on transpiration and evaporation. Transactions American Geophysical Union 25(10).

Toro-Trujillo, A., Ramírez, R., Peña, A., Castillo, A., 2015: Estimation models for the reference evapotranspiration. Agrociencia 49:821–836. https://www.redalyc.org/pdf/302/30243055001.pdf

Trajkovic, S., Kolakovic, S., 2009a: Wind-adjusted Turc equation for estimating reference evapotranspiration at humid European locations. Hydrology Research 40(1), 45–52. https://doi.org/10.2166/nh.2009.002

Trajkovic, S., Kolakovic, S., 2009b: Evaluation of reference evapotranspiration equations under humid conditions. Water Resources Management 23(14):3057–3067. https://doi.org/10.1007/s11269-009-9423-4

Traore, S., Wang, Y. M., Kerh, T., 2008: Modeling reference evapotranspiration by generalized regression neural network in semiarid zone of Africa. WSEAS Transactions on Information Science and Applications 5(6):991–1000.

https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.535.9934&rep=rep1&type=pdf

Tukimat, N. N. A., Harun, S., Shahid, S., 2012: Comparison of different methods in estimating potential<br/>évapotranspiration at Muda Irrigation Scheme of Malaysia. Journal of Agriculture and Rural Development<br/>in the Tropics and Subtropics 113(1), 77–85.<br/>https://www.jarts.info/index.php/jarts/article/view/2012091441739/175

Turc, L., 1961: Evaluation of irrigation water requirements, potential evapotranspiration: A simple climatic formula evolved up to date. Ann. Agron 12(36).

United Nations, 2018: The 2030 Agenda and the Sustainable Development Goals: An opportunity for Latin America and the Caribbean. Goals, Targets and Global Indicators. https://repositorio.cepal.org/bitstream/handle/11362/40156/25/S1801140\_en.pdf, accessed on 21/02/2022.

Urrea, V., Ochoa, A., Mesa, O., 2016: Validación de La Base de Datos de Precipitación CHIRPS Para Colombia a Escala Diaria, Mensual y Anual En El Periodo 1981–2014. XXVII Congreso Latinoamericano de Hidráulica; IAHS, Lima, Perú, accessed on 17.11.2022.

Valipour, M., 2015: Temperature analysis of reference evapotranspiration models. Meteorological Applications 2015:22(3):385–394. https://doi.org/10.1002/met.1465

Wang, K., Dickinson, R. E., 2012: A review of global terrestrial evapotranspiration: observation, modelling, climatology and climatic variability. Reviews of Geophysics 50(RG2005):1–54. https://doi.org/10.1029/2011RG000373

Yang, Z., Liu, Q., Cui, B., 2011: Spatial distribution and temporal variation of reference evapotranspiration during 1961–2006 in the Yellow River Basin, China. Hydrological Sciences Journal 56(6):1015–1026. https://doi.org/10.1080/02626667.2011.590810

Yao, H., Scott, L., Guay, C., Dillon, P., 2009: Hydrological impacts of climate change predicted for an inland lake catchment in Ontario by using monthly water balance analyses Huaxia. Hydrological Processes 23:2368–2382. https://doi.org/DOI: 10.1002/hyp.7347

Yirga, S., 2019: Modelling reference evapotranspiration for Megecha catchment by multiple linear regression. Modeling Earth Systems and Environment 5(2):471–477. https://doi.org/10.1007/s40808-019-00574-2

Yoder, R. E., Odhiambo, L. O., Wright, W. C., 2005: Evaluation of methods for estimating daily reference crop evapotranspiration at a site in the humid southeast United States. Applied Engineering in Agriculture 21(2):197–202. https://doi.org/10.13031/2013.18153

Zanetti, S. S., Sousa, E. F., Carvalho, D. F. de, Bernardo, S., 2008: Estimação da evapotranspiração de referencia no estado do Rio de Janeiro usando redes neurais artificiais. Revista Brasileira de Engenharia Agrícola e Ambiental 12(2):174–180. https://doi.org/10.1590/s1415-43662008000200010

## Meteoro logica



Figure 1. The geographical location of the study area and distribution of climatological stations, principal sinoptic station includes wind velocity data





Figure 2. Methodological diagram of the research



Figure 3. Graphical comparison of evapotranspiration by different methods: a) Aeropuerto Antonio Nariño 1983-2017, b) Obonuco 1983-2017, and c) El Encano 1983-2017



Figure 4. Graphical comparison of machine learning models: a) Aeropuerto Antonio Nariño, b) Obonuco and c) El Encano

Meteoro logica



Figure 5 Estimation of ETo by ANN-M1 at stations with scarce information





Figure 6 Spatialization of ETo in southwestern Colombia

# Meteoro logica

## Artículo en edición

Table 1. Reference evapotranspiration models were used in this research. The input variables for this model are: Ra = extraterrestrial solar radiation (MJ m<sup>-2</sup>), Tmax = maximum temperature (°C) at 2 meters, Tmin = minimum temperature (°C) at 2 meters, Tmean = mean temperature (°C) at 2 meters, Ra = incident solar radiation on the atmosphere (MJ m<sup>-2</sup> d<sup>-1</sup>), Rn = net radiation (MJ m<sup>-2</sup> d<sup>-1</sup>), Rs = net radiation (MJ m<sup>-2</sup>), h= masl, RH = relative humidity (%),  $\Delta$  the slope of the vapor pressure curve [kPa °C<sup>-1</sup>], G = ground heat flux density [MJ m<sup>-2</sup> day<sup>-1</sup>],  $\gamma$  = psychrometric constant [kPa °C<sup>-1</sup>] and U2 the wind speed at 2 m height [m s<sup>-1</sup>]

Name	ETo estimation methods	Reference	Model Type
FAO-PM56	$ETo = \frac{0.408  \Delta(Rn - G) + (\gamma  \frac{900}{Tmean + 273}) U2  (es - ea)}{\Delta + \gamma (1 + 0.34 U2)}$	(Allen et al., 1998)	Combined
Hargreaves	$ET_0 = 0.0023 (Tmean + 17.8) (Tmax - Tmin)^{0.5} Ra$	(Hargreaves and Samani, 1985)	Temperature
Turc	ETo = 0.013 $\left(\frac{1162\pi}{1162}\right)(23.8846xRs + 50)(1 + \frac{50-116}{70})$ if RH< 50	(Turc, 1961)	
	ETo = 0.013 ( $\frac{Tmean}{Tmean+15}$ )(23.8846 <i>Rs</i> + 50) if RH > 50		Radiation
Thornthwaite	$ETo = 16 \ge \left(\frac{10*Tmean}{I}\right)^{\alpha}$ $\alpha = 6.75 \times 10^{-7} \times I^{3} - 7.71 \times 10^{-5} \times I^{2} + 1.79 \times 10^{-2} \times I + 0.49$ $I = \sum_{l=1}^{12} \left(\frac{Tmean}{5}\right)$	(Thornthwaite and Wilm, 1948)	Temperature
Cenicafé	ETo =4.37 <sup>-0,0002*h</sup>	(Jaramillo, 2006)	Temperature



	n performance metrics used: N = Number of data, ETo = observed value FAO-PM56, ETo'
= predicted ETo, $\overline{ET_0}$ = average estimated ETo FAO-PM56, $\overline{ET_0}'$ = average predicted ETo	ted ETo, $\overline{ET_0}$ = average estimated ETo FAO-PM56, $\overline{ET_0}'$ = average predicted ETo

= predicted E10, $E1_0$ = average estimated E10 <i>FAO-PM50</i> , $E1_0$ = average predicted E10							
Name	Equation	Purpose of metrics	Perfect score				
RMSE (Root mean squared error)	$RMSE = \sqrt{\frac{1}{N}\sum_{i=1}^{N}(ETo - ETo')^2}$	Goodness-of-fit for high values	0				
MAE (Mean Absolute Error)	$MAE = \frac{1}{N} \sum_{i=1}^{N}  ETo - ETo' $	Goodness-of-fit for mean values	0				
MBE (Mean Bias Error)	$MBE = \frac{1}{N} \sum_{i=1}^{N} (ETo - ETo')$	Determine the average model bias	0				
NSE (Nash–Sutcliffe efficacy coefficient)	$SE = 1 - \frac{\sum_{i=1}^{N} (ETo - \overline{ET_0})^2}{\sum_{i=1}^{N} (ETo' - \overline{ET_0})^2}$	Evaluate the predictive capability of hydrological models.	1				
R (Pearson correlation coefficient)	$r = \frac{\sum_{i=1}^{N} (ETo - \overline{ET_{0}}) (ETo' - \overline{ET_{0}}')}{\sqrt{\sum_{i=1}^{N} (ETo - \overline{ET_{0}})^{2} * \sum_{i=1}^{N} (ETo' - \overline{ET_{0}}')^{2}}}$	Statistical correlation between two variables	1				

Meteoro logica

## Artículo en edición

Name	Height	Multiy	Multiyear monthly averages of climatological variables				Annual precipitation
Station	masl	Tmax (°C)	Tmed (°C)	Tmin (°C)	RH (%)	SBH (h)	mm year <sup>-1</sup>
Aeropuerto Antonio Nariño	1796	27.3	19.2	12.9	76.5	5.4	1216
Barbacoas	32	32.5	26.2	20	89.5	2.9	6927
Bombona	1493	28.9	20.1	13.9	77.9	4.7	1071
Monopamba	1776	23.6	17.0	11.6	88.8	2.1	3214
Obonuco	2710	20.0	13.0	6.9	79.9	3.3	888
Sindagua	2800	20.4	13.1	7.5	80.2	3.9	988
Botana	2820	19.8	12.6	5.9	78.6	3.2	964
El Encano	2830	18.8	11.7	4.8	86.4	2.5	1402
Taminango	1875	26.2	18.0	13.2	83.9	-	1715

#### Table 3. Characteristics of weather stations in the study area: Tmax, Tmed, Tmin, RH, and SBH

Table 4.1 enormance metres for E16 estimation methods concerning PAO-1 W50							
Performance metrics	Cenicafé	Hargreaves	Thornthwaite	Turc			
RMSE (mm month <sup>-1</sup> )	30.4	20.0	50.4	16.9			
MAE (mm month <sup>-1</sup> )	26.4	18.0	48.7	14.2			
MBE (mm month <sup>-1</sup> )	26.4	-17.9	48.7	10.3			
R	0.80	0.90	0.86	0.81			
NSE	0.64	0.81	0.74	0.65			

Table 4. Performance metrics for ETo estimation methods concerning FAO-PM56

#### Table 5. VIF for climatic variables

Model	Inputs variables	VIF					
		TMAX	TMIN	TMED	RH	SBH	COTA
1	TMAX + TMED	12.59	-	12.59	-	-	-
2	TMAX + SBH	3.07	-	-	-	3.07	-
3	TMAX + SBH + COTA	8.67	-	-	-	3.18	7.93
4	TMAX + SBH + COTA + TMED	12.70	-	32.01	-	3.52	17.89
5	TMAX + SBH + COTA + TMED + RH	12.71	-	41.1	2.63	4.32	25.58
6	TMAX + SBH + COTA + TMED + RH + TMIN	13.29	10.59	51.7	2.64	4.64	26.76



	bold are the best							
Combination	Input parameters	RMSE (mm month <sup>-1</sup> )	MAE (mm month <sup>-1</sup> )	MBE (mm month <sup>-1</sup> )	NSE	R	Observations	
MLR-M1	TMAX, SBH, and COTA	7.7	6.1	0.3	0.86	0.93	All models comply with the error assumptions, and COTA was added as a categorical variable; for MLR-M1,	
MLR-M2	TMAX, SBH, and COTA	5.5	4.4	0.1	0.93	0.93	a natural root transformation was applied, and for MLR- M2 and MLR-M3, a square root transformation was	
MLR-M3	TMAX, HUMREL, and COTA	6.5	5.2	0.1	0.90	0.95	applied.	
ANN-M1	TMAX and COTA	7.3	5.9	0.1	0.87	0.94	In the development of these models, multiple	
ANN-M2	TMAX and TMED	8.2	6.6	0.1	0.84	0.92	architectures were built. The tanh activation function, the	
ANN-M3	TMAX, TMED, and	7.6	6.1	0.1	0.87	0.93	lbgfs optimizer, and a hidden layer with three neurons, a	
	TMIN						learning rate of 0.01 and 1000 iterations were performed for each model. The weights of the COTA with the first, second and third neuron was: 0.010, 0.19 and 0.46 respectively. The weights of the TMAX with the first, second and third neuron was: -0.15, 0.29 and -6.16 respectively. The weights of the hidden layer to the output neuron were: -6.52, 7.62 and -0.30.	
ARIMA-M1	TMAX and COTA	5.2	4.0	-0.1	0.94	0.97	The different combinations between the autoregressive and moving average components with their respective lags were tested. It was obtained that the best combination	
ARIMA-M2	TMAX, TMED, and COTA	5.0	3.9	0.1	0.94	0.97	involved an autoregressive coefficient in the regular and seasonal phases with a seasonality of 12 months. ARIMA $(1,0,0)(1,0,0)_{12}$	
ARIMA-M3	PCA1*COTA + PCA2*COTA	4.1	3.2	-0.1	0.96	0.98		

Table 6. Performance metrics of machine learning models compared to FAO-PM56; the models selected in